

Lecture 3: Climate Change Review

Climate Change Ecology
Geography 404
Jeff Hicke

Radiation budget of Earth




FIGURE 2.9 Earth's Radiation Balance. Approximately 342W/m² of solar energy reaches the Earth's surface. 107W/m² is reflected into space, whereas 235W/m² is emitted from the Earth as long-wave radiation. From Climate Change 2007: The Physical Science Basis. Working Group I Contribution to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.

Hannah, Climate Change Biology, 2011

Global temperature has changed substantially over hundreds of millions of years








FIGURE 2.5 Global Temperature During the Past 500 Million Years. Global mean temperature has fluctuated between icehouse and hothouse conditions during the past billion years. Four major hothouse periods have seen a largely ice-free planet, whereas four major icehouse periods have had major polar or continental ice sheets. Current climate is in a warm phase within an icehouse period. Adapted with permission from Christopher R. Scotese


Hannah, Climate Change Biology, 2011














Changes in global temperature since 1850




FIGURE 2.19 Historic Rise in Global Mean Temperature.
Units are deviation in degrees Celsius from the reference year 1960. Colored lines represent temperature reconstructions using different methods. Bars indicate values from the instrumental record. From Climate Change 2007: The Physical Science Basis, Working Group I Contribution to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.

Hannah, *Climate Change Biology*, 2011

Unprecedented melt on surface of Greenland Ice Sheet



FIGURE 2.22 Recent Greenland Ice Melting.
Satellite images of the ice sheet have shown a dramatic increase in the latter half of the 20th century. Greenland ice loss increases are linked to the melting of sea ice (e.g., in Antarctica), which also has increased in extent because the sea is already melting faster. Continued acceleration of melting could result in a decrease of transatlantic circulation. Source: AVIC Climate Impact Assessment.

Hannah, *Climate Change Biology*, 2011

Changes in atmospheric carbon dioxide






FIGURE 2.16 Mauna Loa CO₂ Record.
The CO₂ record from Mauna Loa clearly shows strongly rising atmospheric CO₂ concentrations during approximately the past 50 years. Superimposed on a multidecadal increase is a much smaller “sawtooth” annual cycle, which results from the release and uptake of CO₂ from vegetation. From Climate Change 2007: The Physical Science Basis, Working Group I Contribution to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.

Hannah, *Climate Change Biology*, 2011


Recent climate change and attribution to humans



Attribution of recent climate change to humans



Climate modeling: CO2 emissions scenarios



Projections of future global temperature





FIGURE 2.29 Global Mean Temperature Estimated from GCMs. Using IPCC emissions scenarios, GCMs simulate global climate change. One summary statistic from these simulations is global mean temperature, shown here as it varies with emissions scenario (colored lines) and GCM (variation around colored lines). From *Climate Change 2007: The Physical Science Basis. Working Group I Contribution to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.*

Hannah, *Climate Change Biology*, 2011

Finer resolution => better studies of impacts



Hannah, *Climate Change Biology*, 2011

Climate modeling: spatial resolution





FIGURE 2.27 RCM Resolution. An RCM can resolve features such as mountain ranges that have important influence on climate. In this example from North America, all mountains from the Sierras of California to the Rocky Mountains are represented as a single "bump" at the horizontal resolution of a GCM, whereas they are better resolved at the resolution of the RCM. *Reproduced with permission from Yale University Press.*

Hannah, *Climate Change Biology*, 2011

Impacts to physical climate system important to biology


Changes in Snowfall Contributions to Wintertime Precipitation 1949 to 2005



Climate Change Ecology 19 Prof. J. Hicke

Impacts to physical climate system important to biology


Observed Spring Snowmelt Dates



Climate Change Ecology 20 Prof. J. Hicke

Impacts to physical climate system important to biology

Trends in Peak Streamflow Timing



Climate Change Ecology 21 Prof. J. Hicke

Impacts to physical climate system important to biology



FIGURE 8.1 Boulder Slides, Queen National Park, 1952. Reproduced with permission from Archives and Special Collections, Mountain Library, The University of British Columbia.



FIGURE 8.2 Boulder Slides, 1982. Reproduced with permission from Archives and Special Collections, Mountain Library, The University of British Columbia.


Hannah, *Climate Change Biology*, 2011

Climate Change Ecology

22

Prof. J. Hicke

Impacts to physical climate system important to biology



Observed changes in relative sea level from 1958 to 2008 for locations on the U.S. coast. Some areas along the Atlantic and Gulf coasts saw increases greater than 8 inches over the past 50 years.


USGCRP, *Global Climate Change Impacts in the United States*, 2009

Climate Change Ecology

23

Prof. J. Hicke

Impacts to physical climate system important to biology



Past and projected sea level rise

FAQ 5.1, Figure 1. Time series of global mean sea level rise relative to the 1992-1999 mean (in the past and as projected for the future). For the period before 1870, global measurements of sea level are not available. The grey shading shows the uncertainty in the estimated long-term rate of sea level change (Section 4.4.3). The red line is a reconstruction of global mean sea level from tide gauges (Section 5.5.2.1), and the red shading shows the range of variations from a smooth curve. The green line shows global mean sea level observed from satellite altimetry. The blue shading represents the range of model projections for the IPCC A1.2 scenario for the 21st century, relative to the 1992 to 1999 mean, and has been calculated independently from the observations. Beyond 2100, the projections are increasingly dependent on the emissions scenario (see Chapter 10 for a discussion of sea level rise projections for other scenarios considered in this report). Over many centuries or millennia, sea level could rise by several meters (Section 10.7.4).


IPCC Working Group I, 2007


Climate Change Ecology


24

Prof. J. Hicke

Impacts to physical climate system important to biology









Hurricanes and warming

Observed Relationship Between Sea Surface Temperatures and Hurricane Power in the North Atlantic Ocean



Observed sea surface temperature (blue) and the Power Dissipation Index (green), which combines frequency, intensity and duration for North Atlantic hurricanes.¹²⁹ Hurricane rainfall and wind speeds are likely to increase in response to human-caused warming. Analyses of model simulations suggest that for each 1.8°F increase in tropical sea surface temperatures, rainfall rates will increase by 6 to 18 percent.¹⁸

Observed and Projected Sea Surface Temperature Change Atlantic Hurricane Formation Region



Observed (black) and projected temperatures (blue = lower scenario, red = higher scenario) in the Atlantic hurricane formation region. Increased intensity of hurricanes is linked to rising sea surface temperatures in the region of the ocean where hurricanes form. The shaded areas show the likely ranges while the lines show the central projections from a set of climate models.
